

***Dacus (Didacus) yemenensis* White, 2006**

Dacus (Didacus) yemenensis White, 2006: 90

Wing length, 7.4 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow dark spot. Frons, frontal setae 2, orbital seta 1 (all very reduced).

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to katepisternum; extended onto katepisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; deep, extending to vein R_{4+5} before wing apex; apically expanded into a spot which reaches vein M and starts before end of vein R_{2+3} .

Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (not as deeply as costal band). Crossbanding; crossbands on R-M and DM-Cu; marking on DM-Cu continued anteriorly to vein R_{4+5} .

Legs. Femora bicoloured (pale basal third to half, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. Aculeus narrowly pointed; no torsion.

(Description after White, 2006)