

***Dacus (Leptoxyda) attenuatus* Collart**

Dacus (Didacus) attenuatus Collart, 1935a: 36

Wing length, 6.2 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot on a dark face. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus and xanthine form uncertain (discoloured); lateral and medial postsutural vitta apparently absent. Scutellum concolorous with scutum (but possibly discoloured). Anepisternal stripe hardly differentiated from general colour of pleura. Laterotergal xanthine confined to katatergite (but barely differentiated from general colour of pleura). Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, reaching at least vein R₄₊₅; apically expanded into a spot which reaches below vein M and starts before crossvein R-M. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. fore- and hindfemora bicoloured (pale basally, fuscous apically); midfemur fuscous.

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)