

***Dacus (Leptoxyda) carnesi* (Munro)**

*Ectopodacus carnesi* Munro, 1984: 85

Wing length, 5.3 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral postsutural vitta absent; medial postsutural vittae present.

Scutellum without any dark patterning (except for basal dark margin).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; fairly deep, extending below vein  $R_{2+3}$  or to vein  $R_{4+5}$ , before wing apex; expanded into a small spot at apex which starts before end of vein  $R_{2+3}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c slightly coloured (not as deeply as costal band). Without any crossbanding.

Legs. Femora pale coloured.

Abdomen. Predominantly fuscous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)