

***Dacus (Leptoxyda) gypsoides* Munro**

Dacus gypsoides Munro, 1933: 4

Wing length, 6.0-7.2 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot on a dark face. Frons, frontal setae 2, orbital setae 0.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin); apical margin concave. Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an almost complete (>90%) covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein R_{4+5} before wing apex; apically expanded into a spot which reaches about mid-depth of cell r_{4+5} and starts before end of vein R_{2+3} . Anal streak present (colour extending beyond cell bcu). Cells bc and c slightly coloured (as deeply as costal band). Without any crossbanding.

Legs. Femora pale coloured.

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)