

***Dacus (Leptoxyda) kakamega* White, 2006**

Dacus (Leptoxyda) kakamega White, 2006: 103

Wing length, 5.4 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral postsutural vitta absent; medial uncertain (position damaged but paler midline area suggests there may be a vitta). Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly shallow, slightly extending below vein R_{2+3} before wing apex (pale below vein); expanded into a small spot at apex. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline.

Crossbanding; a trace of mark at upper end of R-M.

Legs. Femora bicoloured (forefemur pale basal third, mid- and hindfemora pale basal two-thirds, red-brown apically).

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Foretarsomeres 2-5 reddish, contrastingly darker than foretarsomere 1, which is pale (yellowish).

Female. Unknown.

(Description after White, 2006)