

***Dacus (Leptoxyda) masaicus* Munro stat.rev.**

Dacus masaicus Munro, 1937a: 42

Wing length, 9.0-10.5 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine wedge shaped and joined to notopleural callus (but can be reduced); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep); sometimes concolorous with scutum. Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an almost complete (>90%) covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to (or below) vein R_{4+5} before wing apex; apically expanded into a spot which reaches below vein M and starts before end of vein R_{2+3} . Anal streak present (colour extending beyond cell bcu). Cells bc and c slightly coloured (as deeply as costal band). Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data.

(Description after White, 2006)