

***Dacus (Leptoxyda) mediovittatus* White, 2006**

*Dacus (Leptoxyda) mediovittatus* White, 2006: 104

Wing length, 5.0-5.4 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral postsutural vitta absent; medial postsutural vitta present (difficult to discern in MRAC paratype). Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; fairly shallow, slightly extending below vein  $R_{2+3}$  before wing apex; apically expanded into a spot which reaches about mid-depth of cell  $r_{4+5}$  and appears to start before end of vein  $R_{2+3}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (pale basal half to two-thirds, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Foretarsomeres 2-5 reddish, contrastingly darker than foretarsomere 1, which is pale (yellowish).

Female. Unknown.

(Description after White, 2006)