

***Dacus (Leptoxyda) nigrolateris* White, 2006**

*Dacus (Leptoxyda) nigrolateris* White, 2006: 104

Wing length, 7.5 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1 (doubled on left side).

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral postsutural vitta absent; medial postsutural vitta present. Scutellum without any dark patterning (except for deep basal dark margin). Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga without a xanthine, or at most with a trace of a small xanthine on the katatergite (varies from one side of specimen to the other). Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; very deep, extending below vein  $R_{4+5}$  before wing apex; apically expanded into a spot which reaches about mid-depth of cell  $r_{4+5}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Without any crossbanding.

Legs. Forefemur red-brown; midfemur largely red-brown (pale basal quarter, red-brown apically); hindfemur bicoloured (pale slightly more than basal half, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. No available data.

(Description after White, 2006)