

***Dacus (Leptoxyda) parvimaculatus* White, 2006**

*Dacus (Leptoxyda) parvimaculatus* White, 2006: 105

Wing length, 6.4 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a minute dark spot; spot less than half width if 1<sup>st</sup> flagellomere. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral postsutural vitta absent; medial postsutural vitta present. Scutellum without any dark patterning (except for very narrow basal dark margin).

Anepisternum with a narrow stripe from notopleural callus to katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite.

Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Costal band complete; deep, extending to vein R<sub>4+5</sub> before wing apex; apically expanded into a spot (depth uncertain due to damage), and starts before end of vein R<sub>2+3</sub>. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Fore- and midfemora red-brown (somewhat streaked); hindfemur bicoloured (pale basal two-thirds, red-brown apically).

Abdomen. Predominantly fuscous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. No available data.

(Description after White, 2006)