

***Dacus (Psilodacus) externellus* (Munro)**

*Athlodacus externellus* Munro, 1984: 150

Wing length, 5.5 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area dark. Frons, frontal setae 2, orbital seta 1. Thorax. Scutum predominantly black; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine absent; lateral and medial postsutural vittae absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; stripe fairly broad (anteriorly extending to about level of anterior notopleural seta); extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly deep, extending below vein  $R_{2+3}$  or to vein  $R_{4+5}$ , before wing apex; apically expanded into a spot which reaches below vein M and starts before end of vein  $R_{2+3}$ . Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (foreleg only preapically) (pale basally, red-brown apically).

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III without pecten.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)