

***Dacus (Leptoxyda) longistylus* Wiedemann**

Dacus longistylus Wiedemann, 1830: 522

Wing length, 5.2-7.3 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral postsutural vitta usually absent, sometimes present; medial vitta usually present. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep, and may be in the form of a very shallow triangle). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; stripe fairly broad (anteriorly extending to about level of anterior notopleural seta); extended onto katepisternum. Laterotergal xanthine usually absent; sometimes with a xanthine on anatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent (very rarely present). Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with a variable covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent or barely coloured beyond sc. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (preapically or apically) (pale basally, red-brown apically).

Abdomen. Predominantly fulvous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Oviscape about as long as abdomen; aculeus pointed; no torsion.

(Description after White, 2006)