

***Ceratitis (Ceratalaspis) guttiformis* Munro**

Ceratitis (Pardalaspis) guttiformis Munro, 1935: 309.

Body length: 5.80 (5.75-5.85) mm; wing length: 5.03 (4.65-5.40) mm.

Female

Head. Antenna yellow-orange. Third antennal segment three times as long as second segment. Arista with longer hairs but not distinctly plumose. Frons with short scattered hairs which are distinctly darker than frons (in paratype only slightly darker however); concave to flat, in lateral view not distinctly projecting forwards at antennal implant. Chaetotaxy normal for subgenus.

Thorax. Postpronotum concolorous with ground colour of mesonotum, with black spot. Ground colour of mesonotum pale with orange tinge. Chaetotaxy normal for subgenus. Scapular setae black. One anepisternal bristle. Scutellum white basally, otherwise yellow with three black separate markings apically; basally with 2 dark spots (poorly developed in holotype). Subscutellum pale with three dark separate spots.

Legs yellow; setation typical for subgenus, mainly pale especially on femora. Posterior and posterodorsal rows on fore femur pale.

Wing bands with markings extensively yellow. Banding, marginal band continuous; cubital band free or very weakly connected with discal band; medial band absent; crossvein r-m at or before middle of discal cell. Crossvein dm-cu posteriorly more outwards than anteriorly.

Abdomen. Yellow-orange colour. Setation and banding typical for subgenus. Oviscape almost as long as abdominal terga 3-6 combined.

Male

Unknown.

(Description after De Meyer, 1998)