

***Dacus (Leptoxyda) temnopterus* Bezzi**

Dacus temnopterus Bezzi, 1928: 331

Wing length, 5.3 mm.

Head. Pedicel+first flagellomere length uncertain (broken). Face, antennal furrow with a dark stripe. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum red-brown, patterned black; postpronotal lobe concolorous with red-brown areas of scutum; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum with a deep basal band or shallow basal red-brown triangle. Anepisternal stripe exceptionally narrow; not extended onto katepisternum. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an almost complete (>90%) covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; very deep, extending below vein R_{4+5} before wing apex; apically expanded into a spot which reaches about mid-depth of cell r_{4+5} . Anal streak absent (at most with a trace of colour confined to cell bcu). Cell bc hyaline; cell c coloured (as deep as costal band). Crossbanding, absent but there is an expansion of the costal band across R-M.

Legs. Femora red-brown.

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)