

***Dacus (Lophodacus) umehi* White, 2006**

*Dacus (Lophodacus) umehi* White, 2006: 122

Wing length, 4.1 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot continuous with dark face. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral postsutural vitta absent; medial postsutural vitta present. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to katepisternum; stripe fairly broad (anteriorly extending to about level of anterior notopleural seta); extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta absent; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; fairly shallow, slightly extending below vein  $R_{2+3}$  before wing apex; apically expanded into a spot which reaches about mid-depth of cell  $r_{4+5}$  and starts before end of vein  $R_{2+3}$ . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (foreleg only streaked preapically; midfemur pale basal third; hindfemur pale basal two-thirds; red-brown apically).

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Unknown.

(Description after White, 2006)