

***Dacus (Psilodacus) basifasciatus* (Hering)**

Didacus basifasciatus Hering, 1941b: 8

Wing length, 6.6-7.0 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area black. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe bicoloured; notopleural xanthine absent; lateral and medial postsutural vitta absent.

Scutellum without any dark patterning (except for basal dark margin).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; stripe broad (anteriorly extending to before level of anterior notopleural seta); not extended onto katepisternum. Lateroterga without a xanthine. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with microtrichia in a narrow line along centre; remainder bare. Crossvein R-M beyond middle of cell dm. Costal band absent (reduced to a small apical spot). Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Crossbanding; crossband on bm-cu; broad crossband on R-M; mark at upper end of DM-Cu.

Legs. Femora bicoloured (pale basally, fuscous to black apically); forefemur streaked pale.

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. No available data.

(Description after White, 2006)