

***Dacus (Psilodacus) namibiensis* Hancock & Drew**

Dacus (Psilodacus) namibiensis Hancock & Drew, 2001: 74

Wing length, 4.3-4.4 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine absent; lateral and medial postsutural vitta absent.

Scutellum without any dark patterning (except for basal dark margin).

Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; stripe fairly broad (anteriorly extending to about level of anterior notopleural seta); extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with microtrichia confined to anteriorly half. Crossvein R-M beyond middle of cell dm. Costal band reduced to a very narrow band, often absent or barely discernable, and a small apical spot. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora pale coloured.

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)