

***Dacus (Lophodacus) triater* Munro**

Dacus (Psilodacus) triater Munro, 1937b: 2

Wing length, 4.2-5.3 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus concolorous with scutum; notopleural xanthine absent; lateral and medial postsutural vitta absent. Scutellum with a broad black stripe from base to apex (lateral margin yellow); apical margin sometimes concave. Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; stripe broad (anteriorly extending to before level of anterior notopleural seta); not extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br bare or with a few microtrichia, confined to anteriorly half. Crossvein R-M beyond middle of cell dm. Costal band reduced to a very narrow band, often absent or barely discernable, and a small apical spot. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Crossbanding; isolated crossband on R-M.

Legs. Femora pale coloured.

Abdomen. Predominantly black; see image (CD-C). Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)