

***Dacus (Lophodacus) trigonus* Bezzi**

*Dacus trigonus* Bezzi, 1919: 179

Wing length, 5.2-5.3 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, black (possibly comprised of an antennal black spot joined to a dark carina; see image, CD-C). Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine absent; lateral postsutural vitta absent; medial postsutural vitta present. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine very small and confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br without microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent or barely coloured beyond sc. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Crossbanding; a single triangular marking across both R-M and DM-Cu.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly black; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end  $A_1+Cu_2$ , and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. Aculeus pointed and with preapical "shoulder"; no torsion.

(Description after White, 2006)