

***Dacus (Psilodacus) delicatus* Munro**

Dacus (Metidacus) delicatus Munro, 1939a: 31

Wing length, 6.0 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a distinct dark spot (diffuse and very low in HT; more distinct in some specimens). Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe bicoloured; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum with a deep basal band or shallow basal red-brown triangle. Anepisternum with a narrow stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band absent (reduced to a small apical spot). Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (forefemur sometimes only preapically) (pale basally, red-brown apically).

Abdomen. Red-brown, patterned black; shape and patterning, see image.

Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)