

***Dacus (Leptoxyda) fuscovittatus* Graham**

Dacus fuscovittatus Graham, 1910: 169

Wing length, 6.0-6.7 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vittae present (lateral fairly short, barely extending beyond anterior supra-alar seta). Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; fairly shallow, slightly extending below vein R₂₊₃ before wing apex; apically expanded into a spot which reaches below vein M and starts before end of vein R₂₊₃. Anal streak present (colour extending beyond cell bcu). Cells bc and c slightly coloured (not as deeply as costal band). Without any crossbanding.

Legs. Femora bicoloured (pale basally, red-brown apically); hindfemur tending to almost entirely pale.

Abdomen. Predominantly fulvous; shape and patterning, see image. Tergites I-V all fused.

Male. Tergite III with pecten, dense microtrichia adjacent end A₁+Cu₂, and hindtibia preapical "pad". Basal costal sections without specialised setae.

Female. No available data.