

***Dacus (Leptoxyda) opacatus* Munro**

*Dacus opacatus* Munro, 1948b: 19

Wing length, 7.3-7.5 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin, which is deep). Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein R<sub>4+5</sub> before wing apex; apically expanded into a spot which reaches below vein M and starts before crossvein R-M. Anal streak present (colour extending beyond cell bcu). Cell bc hyaline; cell c slightly coloured (not as deep as costal band). Without any crossbanding.

Legs. Forefemur red-brown; mid- and hindfemora bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image. Tergites I-V all fused.

Male. Unknown.

Female. No available data.

(Description after White, 2006)