

***Dacus (Leptoxyda) phantoma* Hering**

Dacus phantoma Hering, 1941b: 6

Wing length, 7.9 mm.

Head. Pedicel+first flagellomere distinctly longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly red-brown; postpronotal lobe concolorous with scutum; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vittae present. Scutellum without any dark patterning (except for basal dark margin, which is sometimes deep). Anepisternum with a narrow stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present.

Wing. Basal cell bc without an almost complete covering of microtrichia; cell c with an almost complete (>90%) covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to vein R_{4+5} before wing apex; apically expanded into a spot which reaches below vein M and starts before crossvein R-M. Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (not as deeply as costal band). Without any crossbanding.

Legs. Fore- and midfemora red-brown; hindfemur bicoloured (pale basally, red-brown apically).

Abdomen. Predominantly red-brown; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Unknown.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)