

***Dacus (Neodacus) xanthaspis* (Munro)**

Tythocalama xanthaspis Munro, 1984: 157

Wing length, 6.7 mm.

Head. Pedicel+first flagellomere slightly longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly fuscous; postpronotal lobe yellow to bicoloured; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum with a deep basal band or shallow basal red-brown triangle. Anepisternum with a narrow stripe from notopleural callus to (or almost to) katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c almost entirely covered in microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; deep, extending to (or below) vein R_{4+5} before wing apex; not expanded into an apical spot. Anal streak present (colour extending beyond cell bcu). Cells bc and c coloured (not as deeply as costal band). Without any crossbanding.

Legs. Femora red-brown.

Abdomen. Predominantly fuscous; shape and patterning, see image (CD-C).

Wasp-waisted, syntergite I+II longer than broad. Tergites I-V all fused.

Male. Unknown.

Female. No available data.

(Description after White, 2006)