

***Ceratitis (Ceratalaspis) lineata* (Hering)**

*Pardalaspis lineata* Hering, 1938: 406.

Body length: 4.60 mm; wing length: 4.50 mm.

Male

(Based on original description, with additions based on KBIN and USNM specimens and indicated as such) Head. Antenna yellow to yellow-orange. Third antennal segment broken in holotype, less than three times as long as second segment in USNM specimen. Arista with short hairs over entire length. Frons (in material studied) yellow, with scattered hairs which are distinctly darker than frons, in lateral view not distinctly projecting forwards at antennal implant; face yellow-white, lower margin near mouth and below eyes more yellow. Chaetotaxy normal for subgenus.

Thorax. Postpronotum yellow-white, spotted. Ground colour of mesonotum pale with orange tinge. Inner scapular setae missing, outer scapular setae black. One anepisternal bristle, in USNM specimen on one side second lower bristle which is less developed. Scutellum white basally, otherwise yellow with three black separate markings apically; basally two dark separate spots. Subscutellum with three black spots, widely separated, dorsally connected.

Wing bands with markings extensively yellow. Banding, marginal band continuous; cubital band free; medial band absent; crossvein r-m at or just before middle of discal cell. Crossvein dm-cu posteriorly more inwards than anteriorly.

Legs yellow; setation typical for subgenus, mainly pale especially on femora. Posterior and posterodorsal rows on fore femur pale. Ventral spines dark in USNM specimen.

Abdomen yellow. Setation and banding typical for subgenus.

Female

Unknown.

(Description after De Meyer, 1998)