

***Dacus (Psilodacus) annulatus* Becker**

Dacus annulatus Becker, 1903: 138

Wing length, 4.9-5.3 mm.

Head. Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot; upper area often dark (especially in males). Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine connected to notopleural callus (but semicircular and believed to be non-homologous with wedge type); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum; anteriorly barely reaching anterior notopleural seta. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; expanded into a small spot at apex. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Pale legged form: femora pale; dark legged form: femora bicoloured (pale basally, apically or preapically fuscous).

Abdomen. Predominantly fuscous; shape and patterning, see image (CD-C).

Tergites I-V all fused.

Male. Tergite III without pecten. Basal costal sections without specialised setae.

Female. Aculeus pointed; no torsion.

(Description after White, 2006)