

***Dacus (Didacus) seguyi* (Munro)**

Pycnodacus seguyi Munro, 1984: 140

Wing length, 5.0 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum predominantly black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum; extended onto katepisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm.

Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; expanded into a small spot at apex. Anal streak absent (at most with a trace of colour confined to cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs. Femora bicoloured (pale basally, mid- and hindfemora red-brown apically; forefemur preapically).

Abdomen. Red-brown, patterned black; shape and patterning, see image (CD-C). Tergites I-V all fused.

Male. Unknown.

Female. Aculeus pointed and with preapical "shoulder"; appears to have torsion.

(Description after White, 2006)