

***Bactrocera (Bactrocera) latfrons* (Hendel)**

Chaetodacus latifrons Hendel, 1915: 425

Wing length, 4.4-6.1 mm.

Head. Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot. Frons, frontal setae 2, orbital seta 1.

Thorax. Scutum black; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine absent; lateral postsutural vitta present; medial postsutural vitta absent. Scutellum without any dark patterning (except for basal dark margin). Anepisternum with a stripe from notopleural callus to (or almost to) katepisternum, extended onto katepisternum; stripe very broad (anteriorly extending to, or almost to postpronotal lobe). Lateroterga with a single xanthine across both anatergite and katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta present; prescutellar acrostichal seta present; basal scutellar seta absent.

Wing. Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not or only slightly extending below vein R_{2+3} before wing apex; expanded into a small apical spot. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding. Legs. Femora pale, sometimes with a dark preapical marking, or bicoloured (pale basally, red-brown to fuscous apically). Abdomen. Red-brown, patterned black. Tergites II-V separate. Male. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical "pad". Basal costal sections without specialised setae. Female. Aculeus pointed and with preapical "shoulder"; no torsion; length, 1.4-1.7mm.

(Description provided by I.M. White)