

***Perilampus woodi*** (Bezzi)

*Carpophthoromyia woodi* Bezzi, 1924: 96.

Body length. 3.40-4.80 mm; wing length 3.85-4.90 mm.

**Male**

Head: Antennal segments brown. Arista short pubescent, longest rays at most equal to width of base of arista. Frons ventral half yellow-white, dorsal part darker yellow. Two frontals, placed parallel to medial eye margin; two orbitals, placed slightly convergent with inner orbital more medially. Face white, below antennal implant with distinct brown band. Occiput in dorsal part with pair of black-brown patches, otherwise dark yellow.

Thorax: Scutum shining brown; dark dispersed pilosity, two transverse bands with silvery pilosity and microtrichosity, one anteriorly of transverse suture, second near dorsocentral setae; second one sometimes missing. Postpronotum white. Anepisternum brown, with white band occupying posterodorsal part, its ventral margin reaching posteroventral corner or almost so; with pale pilosity; one anepisternal seta. Anatergite and katatergite brown. Scutellum white. Subscutellum brown.

Legs: pale yellow, femora and knees brown.

Wing: Anterior part of wing completely brownish coloured by broad band reaching from well below bcu appendix to apex of wing, covering largely cells br and basal half dm, the latter along an oblique line up till where cross-vein R-M touches vein M. Posterior apical band touching former band near middle of cell  $r_{4+5}$ . Basal part of wing completely brownish coloured. R-M ratio 1.92-2.18.

Abdomen: Shining dark black-brown, posterior margin of tergites 2-4 with greyish band, anteriorly more yellow; tergite 5 with median yellow patch.

**Female**

As male. Female terminalia, oviscape at least as long as abdominal tergites, black to black-brown colour, with black pilosity. Aculeus brown, stout cylindrical, about 17 times as long as wide; aculeus tip bluntly pointed, with slightly convex lateral margins, not sinuate.

(Description after De Meyer, 2009)