

***Perilampus dryades* Munro**

Perilampus dryades Munro, 1939: 38.

Body length. 3.75-4.45 mm; wing length 3.80-4.30 mm.

Male

Head: Antennal segments orange. Arista almost bare, at most few dispersed rays shorter than width of base of arista. Frons ventral two-fifths yellow-white, dorsal part orange-brown to brown. Two frontals, placed parallel to medial eye margin; two orbitals, placed slightly convergent with inner orbital more medially. Face white; with brown band or brown patches near antennal implant. Occiput yellow, with two darker, largely confluent, patches in dorsal part.

Thorax: Scutum shining brown; dark dispersed pilosity, median part with silver-grey microtrichosity; two transverse bands with silvery pilosity and microtrichosity, one anteriorly of transverse suture, second band near dorsocentrals. Postpronotum white. Anepisternum brown, with white band occupying posterodorsal part, its ventral margin reaching posteroventral corner or almost so; with pale pilosity; one anepisternal seta. Anatergite and katatergite white. Scutellum white. Subscutellum brown.

Legs: pale yellow, femora yellow-brown.

Wing: Wing bands brown, well developed. Basal part of wing brown, subbasal irregular spots and streaks present. Anterior apical band completely filling cells r_1 and r_{2+3} . Posterior apical band touching anterior apical band. Subapical band isolated. Discal band reaching posterior wing margin; touching anterior apical band near pterostigma; well separated from subbasal spots and streaks, at least in cell cu1. R-M ratio 0.77-0.88.

Abdomen: Shining dark brown, posterior fourth to two-thirds of tergites 2 and 4 with greyish band.

Female

As male. Female terminalia, oviscapae about as long as abdominal tergites, shining orange-brown, with black pilosity. Aculeus flattened, about 10 times longer than wide; apical part pointed with straight sides, sublaterally somewhat broadened; aculeus tip pointed with subapical shoulders.

(Description after De Meyer, 2009)