

***Dacus (Didacus) insolitus* White & Goodger**

Dacus (Didacus) insolitus White & Goodger, 2009: 14.

Size. Very small, wing length, 2.8-3.8 mm.

Male

Head: Pedicel+1st flagellomere not longer than ptilinal suture. Face, antennal furrow without a dark spot. Frons, frontal setae 1, orbital setae 0.

Thorax: Scutum red-brown; postpronotal lobe and notopleural callus concolorous with scutum; notopleural xanthine absent; lateral and medial postsutural vittae absent. Scutellum concolorous with scutum. Anepisternum without a yellow stripe. Katepisternum with a trace of a xanthine. Laterotergal xanthine absent. Setae. Anterior notopleural seta absent; anterior supra-alar seta absent.

Wing: Basal cells bc and c with an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band pale (but distinct) in cell r_1 ; reduced to a very narrow and pale band in cell r_{2+3} (barely discernable); apically expanded into a distinctly coloured oblique spot, reaching vein M. Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Crossbanding; at most with a trace of a crossband on the anterior portion of R-M.

Legs: Femora brown.

Abdomen: Predominantly red-brown, with distinct black pear-drop shaped spots on each of terga III, IV and V, but not coalesced into a strip. Tergites I-V all fused. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical pad.

Female

As male. Aculeus blunt and with preapical "shoulder"; complete torsion.

(Description after White & Goodger, 2009)