

***Dacus (Lophodacus) xanthinus* White & Goodger**

*Dacus (Lophodacus) xanthinus* White & Goodger, 2009: 26.

Size. Small, wing length, 4.2-4.7 mm.

**Male**

Head: Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot, almost a four-sided shape filling width of furrow. Frons, frontal setae 1-2, orbital setae absent.

Thorax: Scutum predominantly red-brown; postpronotal lobe yellow; notopleural callus yellow; notopleural xanthine wedge shaped (connected to notopleural callus), slightly constricted adjacent callus; lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for red-brown basal margin). Anepisternum with a yellow stripe from notopleural callus to katapisternum; extended onto katapisternum. Laterotergal xanthine confined to katatergite. Thoracic setae. Anterior notopleural seta present but reduced (thin); anterior supra-alar seta absent.

Wing: Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br without microtrichia. Crossvein R-M beyond middle of cell dm. Costal band reduced to a very narrow band, often barely discernable, and a small apical spot. Anal streak absent. Cells bc and c hyaline. Without any crossbanding.

Legs: Forefemur pale, mid- and hindfemur pale to barely discernably bicolored (yellow, pale brown apically).

Abdomen: Predominantly red-brown; tergite IV with a round black sublateral spot; tergite V with a broad medial black stripe. Tergites I-V all fused. Tergite III without pecten; no dense microtrichia adjacent end  $A_1+Cu_2$ ; no hindtibia preapical pad. Basal costal section (costagium) with thickened setae.

**Female**

As male. Aculeus retracted and not dissected.

(Description after White & Goodger, 2009)