

***Dacus (Leptoxyda) yaromi* White & Goodger**

Dacus (Leptoxyda) yaromi White & Goodger, 2009: 18.

Size. Small, wing length, 5.6 mm.

Male

Head: Pedicel+first flagellomere not longer than ptilinal suture. Face, antennal furrow with a dark spot; slightly taller than broad, almost filling width of furrow. Frons, frontal setae 2, orbital setae 1.

Thorax: Scutum red-brown; postpronotal lobe concolorous with scutum; notopleural callus red; notopleural xanthine wedge shaped and red (connected to notopleural callus); lateral and medial postsutural vitta absent. Scutellum without any dark patterning (except for basal red-brown margin). Anepisternum with a red subparallel stripe from notopleural callus to katepisternum; extended onto katepisternum. Laterotergal xanthine barely differentiated from red-brown body colour; confined to katatergite. Thoracic setae. Anterior notopleural seta present; anterior supra-alar seta absent.

Wing: Basal cells bc and c without an almost complete covering of microtrichia; cell bm without microtrichia. Narrow subbasal raised section of cell br with extensive covering of microtrichia. Crossvein R-M beyond middle of cell dm. Costal band complete; shallow, not extending below vein R_{2+3} before wing apex; apically expanded into a spot which reaches about mid-depth of cell r_{4+5} ; and starts before end of vein R_{2+3} . Anal streak present (colour extending beyond cell bcu). Cells bc and c hyaline. Without any crossbanding.

Legs: Femora bicoloured (pale basally, brown to black apical third to two-thirds).

Abdomen: Syntergosternite I+II red-brown; terga III-V dark red-brown with a barely differentiated blackish medial stripe; shape similar to *D. attenuatus*. Tergites I-V all fused. Tergite III with pecten, dense microtrichia adjacent end A_1+Cu_2 , and hindtibia preapical pad.

Female

Unknown.

(Description after White & Goodger, 2009)