

***Ceratitis (Ceratalaspis) ovalis* Munro**

Ceratitis (Pardalaspis) ovalis Munro, 1935: 310.

Body length: 4.00 mm; wing length: 4.40 mm.

Male

Head. Antenna yellow-orange. Third antennal segment twice to three times as long as second segment. Arista distinctly plumose, with very long hairs over entire length. Frons with short scattered hairs which are largely the same colour as frons; convex, in lateral view not distinctly projecting forwards at antennal implant. Chaetotaxy normal for subgenus.

Thorax. Postpronotum white, with black spot. Ground colour of mesonotum pale with orange tinge. Chaetotaxy normal for subgenus. Scapular setae obscured in holotype. One anepisternal bristle. Scutellum yellow with three black separate markings apically; basally with two dark spots joining. Subscutellum completely dark.

Legs yellow; setation typical for subgenus, mainly pale especially on femora. Posterior and posterodorsal rows on fore femur pale.

Wing bands with extensive yellow markings. Banding, marginal band continuous; cubital band free; medial band absent; crossvein r-m at middle of discal cell. Crossvein dm-cu posteriorly more inwards than anteriorly. cup extension only weakly sinoid.

Abdomen. Orange-brown colour. Setation and banding typical for subgenus.

Female

Unknown.

(Description after De Meyer, 1998)