

***Ceratitis (Ceratalaspis) sucini* De Meyer**

Ceratitis (Ceratalaspis) sucini De Meyer, 1998 : 287.

Body length: 5.85 mm; wing length: 5.40 mm.

Female

Head. Antenna yellow-orange. Third antennal segment two times as long as second segment. Arista with short hairs over entire length. Frons yellow; with short scattered hairs which are largely the same colour as the frons; in lateral view not distinctly projecting forwards at antennal implant. Chaetotaxy normal for subgenus.

Thorax. Postpronotum white; unspotted. Ground colour of mesonotum pale. Chaetotaxy normal for subgenus. Scapular setae pale. One anepisternal bristle. Scutellum yellow-white; with three black separate markings; basally without distinct spots. Subscutellum pale with three dark separate spots.

Legs yellow; setation typical for subgenus, mainly pale especially on femora. Posterior and posterodorsal rows on fore femur pale. Ventral spines on fore femur reddish.

Wing bands with markings extensively yellow. Banding, marginal band continuous; cubital band free; medial band joined with marginal band; crossvein r-m just before middle of discal cell. Crossvein dm-cu posteriorly more inwards than anteriorly.

Abdomen. Dark yellow. Setation and banding typical for subgenus. Oviscape shorter than abdomen.

Male

Unknown.

(Description after De Meyer, 1998)